

Prion-associated disease

This page was last modified on 18 May 2016, at 10:35.



Contents

- [Condition](#)
- [Individual at risk](#)
- [Guidance at RECRUITMENT for adult volunteer donor and maternal donor \(cord blood donation\)](#)
- [Guidance at CT /WORK-UP](#)
- [Justification for guidance](#)

Condition

Sporadic, familial and variant Creutzfeldt-Jakob disease (CJD), Gerstmann-Straussler-Scheinker diseases and fatal familial insomnia.

Individual at risk

Recipient

Guidance at RECRUITMENT for adult volunteer donor and maternal donor (cord blood donation)

Unacceptable if:

1. diagnosed with any form of CJD or other prion-associated disorder.
2. identified as at increased risk of developing a prion-associated disorder. This includes:
 - a) individuals at familial risk of developing a prion-associated disorder (have had two or more blood relatives develop a prion-associated disorder or have been informed they are at risk following genetic counselling.)
 - b) individuals who have been told they have been put at increased risk from surgery, transfusion or transplant of tissues and organs.
 - c) individuals who have been told they may be at increased risk because a recipient of blood or tissues they have donated has developed a prion-related disorder.
 - d) recipients of dura mater grafts
 - e) recipients of corneal, scleral or other ocular tissue grafts.
 - f) recipients of human pituitary derived extracts eg Metrodin, human growth hormone.

Guidance at CT/WORK-UP

List as above.

If the donor is asymptomatic then they may proceed at the discretion of the requesting transplant centre.

Justification for guidance

Prion-associated diseases are transmissible by transfusion of blood and likely to be transmissible by stem cells too.